

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

SCIENCE REFERENCE LIBRARY

11

Publication number

0 130 523

A2

D10 CF

12

EUROPEAN PATENT APPLICATION

21 Application number: 84107266.3

22 Date of filing: 25.06.84

51 Int. Cl.: C 12 Q 1/68
// G01N33/548, G01N33/547

30 Priority: 05.07.83 US 511064
18.05.84 US 611667

43 Date of publication of application: 09.01.85
Bulletin 85/2

64 Designated Contracting States: CH DE FR GB IT LI NL
SE

71 Applicant: Molecular Diagnostics, Inc., 400 Morgan
Lane, West Haven, CT.06516 (US)

72 Inventor: Dattagupta, Nanibhushan, 470 Prospect Street,
New haven, Ct 06511 (US)
Inventor: Crothers, Donald M., Surrey Drive, Northford,
Ct 06472 (US)

74 Representative: Adrian, Albert, Dr. et al, c/o BAYER AG
Konzernverwaltung RP Patentabteilung,
D-5090 Leverkusen 1, Bayerwerk (DE)

54 Immobilized nucleic acid probe and solid support for nucleic acids.

57 A solid support capable of binding a nucleic acid thereto upon suitable irradiation, comprising (a) a solid substrate, (b) a photochemically reactive intercalator compound or other nucleic acid-binding ligands, and (c) divalent radical chemically linking the substrate and the ligand (b). Specifically, a hydroxy group-containing solid substrate such as nitrocellulose paper is linked via a bifunctional reagent such as cyanogen bromide or 1,4-butanediol-diglycidyl ether to an amino-substituted angelicin or psoralen or ethidium bromide which in turn is photochemically linked to a nucleic acid. The resulting immobilized nucleic acid probe is capable of hybridizing with complementary nucleic acid fragments and is thereby useful in diagnostic assays.

EP 0 130 523 A2

5 The present invention relates to a novel way of joining a nucleic acid to a solid substrate so as to provide a solid probe suitable for use in various tests, particularly hybridization assays for the determination of specific polynucleotide sequences.

10 In DNA-DNA hybridization and DNA-RNA hybridization assays, one of the complementary nucleic acid chains is commonly coupled to a solid support. This helps to reduce the background and can be used to separate or isolate the corresponding nucleic acid. The methods of attachment of DNA to a solid support have involved (1) non-specific physical adsorption of a single-stranded
15 DNA to nitrocellulose papers, and (2) covalent attachment via diazo coupling. Both methods are specific for single-stranded DNA. These covalent reactions are non-specific and several sites are coupled. These cause ineffective hybridization and loss
20 of perfect fidelity. Several points of attachment per chain reduces the flexibility of the DNA and reduces the rate of hybridization. Moreover, the lifetime of such an adduct is not very long. The DNA comes off easily and it is difficult to quantify the amount on the solid
25 support, without the use of radioactivity. The use of DNA probes for diagnostic purposes demands an effective method of tagging the DNA to a phase which can be separated easily from the rest of the nucleic acids. U.S. Application Serial No. 511,063, filed on July 5,
30 1983, now pending in the names of Dattagupta, Rae and Crothers discloses various tests for nucleic acids, e.g., DNA of individuals being tested for sickle cell anemia. The test involves a soluble labelled probe and a probe fixed to a solid support. The probe can be
35 fixed to the support chemically as by a bifunctional reagent which at one end reacts with the support, e.g., a hydroxyl group of a cellulose molecule, and at the

other end reacts with the DNA. This is quite satisfactory for many purposes but in some instances there may be too much bonding between the substrate and DNA, impairing the sensitivity of the DNA in the test.

5 It is accordingly an object of the present invention to provide a way of binding a nucleic acid to a solid substrate easily and without impairing the sensitivity of the DNA in the test.

10 These and other objects and advantages are realized in accordance with the present invention wherein there is provided a solid support capable of binding a nucleic acid thereto upon suitable irradiation, comprising (a) a solid substrate, (b) a photochemically reactive nucleic acid-binding ligand, and (c) a divalent radical
15 chemically linking the substrate and the nucleic acid-binding ligand.

The specific coupling reagents employed are functionalized, photochemically reactive nucleic acid-binding ligands, e.g., intercalator compounds such
20 as amino-substituted furocoumarins, e.g., amino-methyl-dimethyl-angelicin and amino-methyl-trimethyl-psoralen, and aminophenanthridium halides as well as closely related chemical derivatives thereof, and non-intercalator compounds such as netropsin, distamycin, Hoechst
25 33258 and bis-benzimidazole. Upon photoactivation these reagents will chemically link with nucleic acids. These reagents have a functionalized site other than the nucleic acid-reactive site and, by such other site, they are joined to a solid substrate, thereby in turn joining
30 the nucleic acid to such substrate with a minimum impairment of the nucleic acid function.

Apparently functionalized and photochemically reactive forms of a wide variety of intercalating agents can be used as the coupling reagent as exemplified in
35 the following table:

Intercalator Classes and
Representative Compounds

Literature References

5	A. Acridine dyes	Lerman, J. Mol. Biol. 3:18(1961); Bloomfield et al, "Physical Chemistry of Nucleic Acids", Chapter 7, pp. 429-476, Harper and Rowe, NY(1974) Miller et al, Bio- polymers 19:2091(1980)
	proflavin, acridine orange, quinacrine, acriflavine	
10	B. Phenanthridines	Bloomfield et al, supra; Miller et al, supra
	ethidium coralyne	Wilson et al, J. Med. Chem. 19:1261(1976)
15	ellipticine, ellipticine cation and derivatives	Festy et al, FEBS Letters 17:321(1971); Kohn et al, Cancer Res. 35:71(1976); LePecq et al, PNAS (USA)71: 5078(1974); Pelaprat et al, J. Med. Chem. 23:1330(1980)
20	C. Phenazines 5-methylphenazine cation	Bloomfield et al, supra
	D. Phenothiazines chlopramazine	ibid
25	E. Quinolines chloroquine quinine	ibid
	F. Aflatoxin	ibid
30	G. Polycyclic hydrocarbons and their oxirane derivatives	ibid
	3,4-benzpyrene, benzpyrene diol epoxide, 1-pyrenyl- oxirane	Yang et al, Biochem. Biophys. Res. Comm. 82:929(1978)
35	benzanthracene-5,6-oxide	Amea et al, Science 176:47(1972)

	H. Actinomycins actinomycin D	Bloomfield et al, supra
	I. Anthracyclines β-rhodomyacin A daunomycin	ibid
5	J. Thiazanthrenones miracil D	ibid
	K. Anthramycin	ibid
10	L. Mitomycin	Ogawa et al, Nucl. Acids Res., Spec. Publ. 3:79(1977); Akhtar et al, Can. J. Chem. 53:2891(1975)
	M. Platinum Complexes	Lippard, Accts. Chem. Res. 11:211(1978)
15	N. Polyintercalators echinomycin	Waring et al, Nature 252:653(1974); Wakelin, Biochem. J. 157:721(1976)
20	quinomycin triostin BBM928A tandem	Lee et al, Biochem. J. 173:115(1978); Huang et al, Biochem. 19: 5537(1980); Viswamitra et al, Nature 289: 817(1981)
25	diacridines	LePecq et al, PNAS (USA)72:2915(1975); Carrellakis et al, Biochim. Biophys. Acta 418:277(1976); Wakelin et al, Biochem 17:5057(1978); Wakelin et al, FEBS Lett. 104:261(1979); Capelle et al, Biochem. 18:3354(1979); Wright et al, Biochem. 19:5825(1980); Bernier et al, Biochem. J. 199:479 (1981); King et al, Biochem. 21:4982(1982)
30		
35	ethidium dimer	Gaugain et al, Biochem. 17:5078(1978); Kuhlman et al, Nucl. Acids Res.

5:2629(1978); Maricovits
et al, Anal. Biochem.
94:259(1979); Dervan et
al, JACS 100:1968(1978);
ibid 101:3664(1979).

- 5 ellipticene dimers
 and analogs Debarre et al, Compt.
 Rend. Ser. D. 284:
 81(1977); Pelaprat et
 al, J. Med. Chem.
 23:1336(1980)
- heterodimers Cain et al, J. Med.
10 Chem. 21:658(1978);
 Gaugain et al, Biochem.
 17:5078(1978)
- trimers Hansen et al, JCS
 Chem. Comm. 162(1983);
 Atnell et al, JACS
15 105:2913(1983)
- O. Norphillin A Loun et al, JACS 104:
 3213(1982)
- P. Fluorenes and fluorenones Bloomfield et al, supra
- fluorenodiamines Witkowski et al,
20 Wiss. Beitr.-Martin-
 Luther-Univ. Halle
 Wittenberg, 11(1981)
- Q. Furocoumarins
- angelicin Venema et al, MGG,
25 Mol. Gen. Genet.
 179;1 (1980)
- 4,5'-dimethylangelicin Vedaldi et al, Chem.-
 Biol. Interact. 36:
 275(1981)
- psoralen Marciani et al, Z.
30 Naturforsch B 27(2):
 196(1972)
- 8-methoxypsoralen Belognzov et al, Mutat.
 Res. 84:11(1981);
 Scott et al, Photochem.
 Photobiol. 34:63(1981)
- 35 5-aminomethyl-8-
 methoxypsoralen Hansen et al, Tet. Lett.
 22:1847(1981)
- 4,5,8-trimethylpsoralen Ben-Hur et al,

Biochim. Biophys.
Acta 331:181(1973)

4'-aminomethyl-4,5,8-
trimethylpsoralen

Issacs et al, Biochem.
16:1058(1977)

5

xanthotoxin

Beaumont et al,
Biochim. Biophys.
Acta 608:1829(1980)

R. Benzodipyrones

Murx et al, J. Het.
Chem. 12:417(1975);
Horter et al, Photo-
chem. Photobiol. 20:
407(1974)


10

S. Monostral Fast Blue

Juarranz et al, Acta
Histochem. 70:130 (1982)

15

Angelicin, more accurately 4'-aminomethyl-4,5'-
dimethylangelicin, has the structural formula




20

25

(see Dall'Acquz et al, Photochemistry and Photobiology,
Vol. 37, No. 4, pp. 373-379, 1983.)

Psoralen, more accurately 4'-aminomethyl-4,5',8-
tri-methyl-psoralen (AMT), has the structural formula

30




35


0130523

(Cadet et al, Photochemistry and Photobiology, Vol. 37,
No. 4, pp. 363-371, 1983.)

Methidium chloride, for example, has the formula



(see Graves et al, Biochemistry, 1981, Vol. 20 pp.
15 1887-1892.) Its mono- and di-azide analogues, shown
below, are comparably reactive:



methidium monoazide

methidium diazide

as are the ethyl counterparts and the 4-(3-aminopropyl-
N-carbamoyl) derivative of the phenyl side chain
30 (methidium propylamine).

The solid substrate can be any solid which has
reactive groups which could be carboxyl, amino or the
like, but the preferred reactive groups are hydroxyl
such as are found on cellulose. The cellulose may be
35 unmodified as in cotton or paper or regenerated as in
rayon or partially esterified as in cellulose acetate,
cellulose propionate and especially cellulose nitrate,

CONFIDENTIAL

0130523

or partially etherified as in methylcellulose and carboxymethylcellulose.

5 While the photochemically active intercalator reagent could be directly combined with the solid substrate, advantageously there is a mutual coupler which makes the connection. Suitable reagents include bifunctional compounds such as cyanogen bromide (CNBr), 1,4-butanediol diglycidyl ether, and the like. These are reacted with both the solid substrate and the
10 photochemical reagent simultaneously or first with one and then with the other.




Thereafter, the product is further reacted with the nucleic acid photochemically. The reactions with the coupler and nucleic acid are substantially
15 quantitative so the quantities of the reagents employed depend upon the desired ratio of nucleic acid to solid support. For most purposes, about 0.1 to 1000 mg and preferably about 1 to 100 mg of nucleic acid per gram of solid support will be suitable, although it may be
20 higher or lower, depending upon the molecular weight of the nucleic acid, its sensitivity and the particular test in which it is to be used.

The reaction conditions in each step are generally known per se and any solvents and temperatures can be
25 employed which permit the reactions to proceed without interference, e.g., from about -10 to 100°C., preferably about 10 to 50°C., and most preferably room temperature, employing inert organic solvents such as ether, carbon tetrachloride, THF, and the like.

30 The photochemically active reagents herein employed preferably react through amino groups. Identifying it as RNH_2 and the substrate with pendent OH groups as S, the stepwise reactions are as follows:

35

A. a) with CNBr



B. $\textcircled{\text{S}}$ ~~~~~ NHR

nucleic acid ~~~~~ irradiation



Amino-derivatives of angelicin and psoralen react correspondingly, if not identically.

The particular wavelength of radiation selected will depend upon the particular photoreagent and whether it is desired to bind to a single strand of nucleic acid or to a double strand. If to both strands it can be in a manner and to a degree such that the nucleic acid is no longer denaturable.

0130523

The nucleic acid can be RNA or DNA or short (oligonucleotide) or long chain length, as desired, doubly or singly stranded.

5 Formation of monoadducts is desirable for hybridization experiments. In crosslinks, both DNA strands are covalently linked to psoralen chromophore and hence strand separation prior to hybridization is difficult. If the probe to be hybridized is linked to another non-specific piece of DNA, the non-specific
10 part can be linked either via crosslink or monoadduct formation. In this case, irradiation can be done at any wavelengths between 300-390 nm. Irradiation at 390 nm produces monoadduct; irradiation at 360-300nm produces both monoadduct and crosslinks.

15 If angelicin compounds are used, the product will predominantly be monoadduct irrespective of the wavelength of irradiation.

The invention will now be further described with reference to the accompanying examples wherein all
20 parts are by weight unless otherwise expressed.

Example:

1. Activation of the solid support and coupling of AMT.

25 The procedure described below has been followed for Sephadex G25 and cellulose, but any hydroxy-containing solid support can be activated by an identical procedure.

30 a) Activation with 1,4-butane-diol-diglycidyl ether.

0.5-1 gm solid powder is swollen with water and washed, then 5-10 ml sodium hydroxide solution (0.5 M) is added. To this thick suspension, 1 ml
35 1,4-butanediol-diglycidyl ether is added. The suspension is shaken overnight on a mechanical shaker and then washed with sodium hydroxide (0.5 M) solution and 1.0 ml 4'-aminomethyl-4,5',8-trimethyl-psoralen (2 mg/ml) in water is added, followed by enough 1 M sodium

hydroxide to have a thick suspension. The suspension is then stirred gently for 24 hours at room temperature and excess unreacted residues are quenched with lysine.

5 The solid is then washed with water followed by the desired aqueous buffer solution for DNA coupling.

b) For epoxidation of paper the identical procedure is followed with Whatman filter papers type 540, 1 and 541. The filter papers are taken on a watch glass or beaker cover (glass) and turned occasionally
10 by hand. The rest of the procedure is the same as above.

c) Activation by cyanogen bromide and coupling of AMT. Typical example with cellulose:

0.5 gm cellulose is swollen in 5.0 ml distilled
15 water for one hour. The swollen gel is washed thoroughly with distilled water. Then it is taken in an erlenmeyer flask, ice-cooled distilled water is added to the swollen cellulose and the pH is adjusted between 10.5 - 11 with 5M sodium hydroxide solution.
20 The flask with its contents is cooled in ice to avoid temperature rise above 15°C. 1 gm of solid cyanogen bromide is added to the cellulose and the solution is stirred for 30 minutes and pH maintained between 10.5 - 11 by NaOH. The suspension is washed with ice cold
25 distilled water, water is removed by centrifugation and 20 ml ice cold potassium phosphate buffer (10 mM; pH 8) is added. The activated cellulose is kept in brown bottles (in small aliquots) at - 20°C.

2 - 3 ml of swollen, activated gel is taken in a
30 brown bottle and 0.7 ml AMT (2mg/ml) is added and the mixture is shaken gently in the cold room. Excess activated residues are quenched with lysine. The solid is washed with aqueous buffer for DNA binding.

d) For papers, similar procedures have been
35 followed with Whatman cellulose filter papers type 540, 1 and 541 quantitative papers. Care should be taken to avoid tearing of the papers.

e) Parallel experiments with ^3H labelled aminomethylpsoralen or angelicin are used to estimate labelling efficiency.

2. Coupling of phenanthridium compounds to a solid support and azide formation for photochemical coupling of DNA:

Activation of the solid supports is done by the method described above. As an example, methidium propylamine (R.P. Hertzberg and P.B. Dervan, JACS, 104, 313 (1982)) is coupled to the solid support, using identical buffer conditions as in 1. The isolated methidium containing solid support is then diazotized and azide derivative is made as follows. 1 gm cellulose or (2x5 cm²) of a sheet of activated paper containing methidium chloride is taken in 20 ml water, cooled in ice, 0.2 ml ice cold HCl is added; sodium azide (20 mg solid; 2x) is added. The vessel is cooled in ice and sodium nitrate solid (100 mg) is added. The reaction is allowed to proceed for 30 minutes, solid support is washed with the desired buffer. Coupling of DNA and hybridization are carried out the same way as described for aminomethyl-psoralen. Aminomethyl-dimethyl-angelicin can be similarly treated.

3. Photochemical coupling of DNA:

0.5 ml (0.2 - 0.3 gm gel + buffer) activated solid powder or 0.8 x 1 cm² activated paper is taken in a 1 cm path length spectrophotometer cuvette. Adenovirus DNA (partially labelled with ^3H) (concentration 25 µg/ml) in tris EDTA buffer (10 mM tris, 1mM EDTA, pH 7.5) is added to the cuvette and irradiation is done at a desired wavelength for 30 minutes to two hours depending on the future needs. For AMT, irradiation at 390 nm produces monoadduct whereas at 360 - 300 nm both monoadduct and crosslinks are formed. By altering the concentration and DNA sequence, crosslink to monoadduct formation can be modulated. After photoirradiation, the solid is washed and the radioactivity of the

washings and the solid support is counted in a Beckman 7800 scintillation counter.

Typical Results

5	<u>Solid support</u>	<u>% Coupling</u>	<u>DNA Coupled µg</u>
	0.5ml or 0.8 x 1 control paper (No DNA)	-	-
10	BDGE treated paper	80	20
	Cellulose cellex CNBr activated	91.5	22.5
15	Cellulose cellex BDGE activated	93.4	22.5
20	Sephadex G25 CNBr activated	69.5	18.0

4. Assay for DNA-DNA hybridization of DNA photochemically coupled to the solid support: Adenovirus DNA is covalently coupled to the solid support as above and hybridization with ³H labelled adenovirus DNA is done following the procedure of Noyes and Stark, Cell, 5, 301-310 (1975).

5. Use of photochemically coupled DNA for sickle cell diagnosis: AMT coupled DNA can be recovered as free DNA by irradiation at 260 nm. The separation probe (Application Serial No. 511,063, supra) is coupled to the solid support by the method described above. Then the support with the coupled DNA is mixed with the unknown and the detection probe under hybridization condition - as in 4. The solid support is then tested for the presence of label. If a radioactively labelled detection probe is used, radioactivity is counted.

5a. The product of 4 is irradiated at 260 nm in otherwise the same manner as in 3, whereupon the DNA uncouples from the solid support, entering the solvent medium, viz. aqueous buffer. Then the liquid is
5 assayed for ^3H .

It will be understood that the specification and examples are illustrative but not limitative of the present invention and that other embodiments within the spirit and scope of the invention will suggest
10 themselves to those skilled in the art.

15

20

25

30

35

CLAIMS:

1. A solid support capable of binding a nucleic acid thereto upon irradiation, comprising (a) a solid substrate, (b) a photochemically reactive nucleic acid-binding ligand, and (c) a divalent radical chemically linking the substrate and the nucleic acid-binding ligand.

2. A support according to claim 1, characterized in that the nucleic acid-binding ligand is an intercalator compound selected from acridine dyes, phenanthridines, phenazines, furocoumarins, phenothiazines, and quinolines.

3. A support according to claim 1 or 2, characterized in that the substrate (a) in free state has free OH groups through which it is linked by the divalent radical (c).

4. A support according to any of claims 1 to 3, characterized in that the substrate (a) is cellulose or a cellulose ester.

5. A support according to any of claims 1 to 4, characterized in that the divalent radical (c) is derived from cyanogen bromide or from 1,4-butanediol-diglycidyl ether.

6. A support according to any of claims 1 to 5, characterized in that (b) is psoralen, angelicin, ethidium or derivatives thereof.

7. An immobilized nucleic acid probe comprising (a) a nucleic acid, (b) a nucleic acid-binding ligand photochemically linked to the nucleic acid, and (c) a

8. An immobilized probe according to claim 7, characterized in that the nucleic acid-binding ligand is an intercalator compound selected from acridine dyes, phenanthridines, phenazines, furocoumarins, phenothiazines, and quinolines.

9. An immobilized probe according to claim 7 or 8, characterized in that the substrate (c) in free state
10 has free OH groups through which it is linked by the
divalent radical.

MD 202

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

SCIENCE REFERENCE LIBRARY

(11) Publication number:

0 130 523
A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 84107266.3

(51) Int. Cl.⁴: **C 12 Q 1/68**
//G01N33/548, G01N33/547

(22) Date of filing: 25.06.84

(30) Priority: 05.07.83 US 511064
18.05.84 US 611667

(43) Date of publication of application:
09.01.85 Bulletin 85/2

(88) Date of deferred publication of search report: 23.07.86

(84) Designated Contracting States:
CH DE FR GB IT LI NL SE

(71) Applicant: Molecular Diagnostics, Inc.
400 Morgan Lane
West Haven, CT.06516(US)

(72) Inventor: Dattagupta, Nanibhushan
470 Prospect Street
New haven, Ct 06511(US)

(72) Inventor: Crothers, Donald M.
Surrey Drive
Northford, Ct 06472(US)

(74) Representative: Adrian, Albert, Dr. et al,
c/o BAYER AG Konzernverwaltung RP Patentabteilung
D-5090 Leverkusen 1, Bayerwerk(DE)

(54) Immobilized nucleic acid probe and solid support for nucleic acids.

(57) A solid support capable of binding a nucleic acid thereto upon suitable irradiation, comprising (a) a solid substrate, (b) a photochemically reactive intercalator compound or other nucleic acid-binding ligands, and (c) divalent radical chemically linking the substrate and the ligand (b). Specifically, a hydroxy group-containing solid substrate such as nitrocellulose paper is linked via a bifunctional reagent such as cyanogen bromide or 1,4-butanedioldiglycidyl ether to an amino-substituted angelicin or psoralen or ethidium bromide which in turn is photochemically linked to a nucleic acid. The resulting immobilized nucleic acid probe is capable of hybridizing with complementary nucleic acid fragments and is thereby useful in diagnostic assays.

EP 0 130 523 A3



European Patent
Office

EUROPEAN SEARCH REPORT

0130523
Application number

EP 84 10 7266

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	CHEMICAL ABSTRACTS, vol. 88, no. 11, 13th March 1978, page 99, no. 69689r, Columbus, Ohio, US; L. KITTLER et al.: "Photochemically induced binding of furocoumarins with lambda phage DNA in situ", & STUD. BIOPHYS. 1977, 66(3), 237-41 * Abstract *	1,2	C 12 Q 1/68 X/ G 01 N 33/548 G 01 N 33/547
A	--- US-A-4 169 204 (J. HEARST et al.) * Abstract; column 1 *	1,2	
A	--- CHEMICAL ABSTRACTS, vol. 86, no. 19, 9th May 1977, page 118, no. 134373h, Columbus, Ohio, US; P. CHANDRA: "Nucleic acid modification by furocoumarins and light: some biomedical implications", & PHOTOCHEMOTHERAPIE, VERHANDLUNGSBER. DTSCH.-SCHWED. SYMP. PHOTOMEDIZIN 1975 (Pub. 1976, 25-32 * Abstract *	1,2	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 12 Q G 01 N
--- The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 23-04-1986	Examiner OSBORNE H.H.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT

Page 2

DOCUMENTS CONSIDERED TO BE RELEVANT			Page 2
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	CHEMICAL ABSTRACTS, vol. 98, no. 9, 28th February 1983, page 315, no. 68377g, Columbus, Ohio, US; H. BUENEMANN: "Immobilization of denatured DNA to macroporous supports : II. Steric and kinetic parameters of heterogeneous hybridization reactions", & NUCLEIC ACIDS RES. 1982, 10(22), 7181-96 * Abstract *	1,2,5	
A	--- CHEMICAL ABSTRACTS, vol. 96, no. 7, 15th February 1982, page 207, no. 47770y, Columbus, Ohio, US; R.P. HERTZBERG: "Cleavage of double helical DNA by methidium-propyl-EDTA-iron(II), & J. AM. CHEM. SOC. 1982, 104(1), 313-15 * Abstract *	1,2	
A	--- PROC. NATL. ACAD. SCI. USA, vol. 79, no. 15, August 1982, pages 4594-4598; W.A. SAFFRAN et al.: "Site-directed psoralen crosslinking of DNA" * Page 4594, abstract; page 4598, discussion, final paragraph * --- -/-	1,2	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
Place of search THE HAGUE		Date of completion of the search 23-04-1986	Examiner OSBORNE H. H.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			Page 3
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
A	CHEMICAL ABSTRACTS, vol. 96, no. 5, 1st February 1982, page 310, no. 31231n, Columbus, Ohio, US; & DD - A - 148 955 (AKADEMIE DER WISSENSCHAFTEN DER DDR, ZENTRALINSTITUT FUER MOLEKULARBIOLOGIE) 17-06-1981 * Abstract *	1-4	
A	EP-A-0 070 687 (STANDARD OIL CO.) * Page 4, paragraph 2; page 6, line 20 - page 7, line 14 *	1-4	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 23-04-1986	Examiner OSBORNE H.H.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	